

Results on new drugs from Survey on tobacco, alcohol and other drugs 2011-2012: Slovenia

Darja Lavtar and Andreja Drev National Institute of Public Health

GPS meeting, EMCDDA, Lisbon, 18 June 2014

Outline

- Data collection and methodology
- New drugs / Slovenia
- Questions in the survey
- Results
 - Life time prevalence of drug use
 - Prevalences, age of onset, new drugs used
 - Life time prevalence of drug use by sociodemographic variables



Data collection and methodology

- Based on EMCDDA Handbook for Surveys on Drug Use among the General Population (2002) and national references and practices NIJZ prepared recommendations for implementation of the survey in Slovenia
- Data collection: 2011 (Oct-Nov) and 2012 (Apr-Jun)
- Target population: inhabitants of Slovenia aged 15–64 years, living in private households
- Mixed mode of interview: web, telephone, face-to-face
- Sample size: 7514 respondents
- Response rate: 52,9 %



New drugs

- New drugs: a broad spectrum of psychoactive substances that newly appeared on the drugs market or their abuse appeared only recently and they are thus not yet listed as illicit drugs.
- In July 2013, the Government of the Republic of Slovenia confirmed the Decree Amending the Decree on the Scheduling of Illicit Drugs with which 48 substances were listed as illicit drugs.
- At the time of the implementation of the Survey on Tobacco, Alcohol and Other Drugs, mephedrone was the only new drug that had already been listed as an illicit drug in Slovenia.



Questions: the use of other (new) drugs

- Asked in the same way as the use of drugs in EMQ:
 - Personally knowing people who take other drugs (spice, mephedron, methylone, etc)
 - Lifetime prevalence
 - Name(s) of drugs used lifetime
 - Age of onset
 - Last year prevalence
 - Last month prevalence
 - Last month frequency (in days)
 - Name(s) of drugs used in the last month



Life time prevalence of drug use






Figure 1: Lifetime prevalence of illicit drug use in the Slovenian population aged 15–64 by type of illicit drug

Use of new drugs

- Prevalence of new drugs use:
 - life time prevalence: 0,6 %
 - last year prevalence: 0,3 %
 - last month prevalence: 0,1 %
- Average age at first new drug use: 21 years old
- Drugs reported in the survey: methylone and mephedrone.
- Current situation (reported by DrogArt association and EWS Slovenia): 3-MMC is most widely used new drug at the moment.



Life time prevalence of new drugs

	Socio-demographic category	LTP
GENDER	Male	0,9%
	Female	0,3%
AGE	15-24 years	1,8%
	25-34 years	1,0%
	35-44 years	0,4%
	45-54 years	0,1%
	55-64 years	0,1%
EDUCATION	Primary school or lower	0,7%
	Secondary school	0,6%
	Tertiary education	0,5%
STATUS	Employed, self-employed, farmer	0,3%
	Primary, secondary or tertiary student	2,0%
	Retired person	0,0%
	Unemployed	1,6%



Thank you for your attention!

Contact:

darja.lavtar@nijz.si

