

STANDARD TABLE 1

STANDARDIZED RESULTS AND METHODOLOGY OF ADULTS NATIONAL POPULATION SURVEYS ON DRUG USE

1. - General Information

1.1. - Notes:

For printing, please use the "Landscape" mode

(EMQ) indicates the substances included in the European Model Questionnaire

For more information on the EMQ, please [see](#) .

For detailed information on the methodology of national surveys, please see the [Statistical Bulletin](#) .

Guidelines regarding substances

When information from a substance is available, please fill in all cells (M,F,T) and for Lifetime; Last 12 months and Last 30 days.

When a substance was not asked for (information not available) leave cells empty in the table, and give that information in "Drug Definitions"

Guidelines to use decimal figures and to fill in cells

- Please provide figures with one decimal place (even if the decimal is 0 (e.g. 15.2; 3.6; 6.0).

- Round off the second decimal place to the nearest first decimal. (e.g. 2.31 -->2.3; 6.35 -->6.4; 0.08 -->0.1)

- When there are not cases in a cell, use 0.0 (*)

- When there is no information -substance not asked for- leave cells empty. See below also

(*) This rule will make equivalent values <0.05% (less than one case in 2000 population) and 0.00%.

The alphanumeric symbols used sometimes to indicate "no cases" in a cell (such as - or *) create problems in the Epidemiological database

"CI 95" stands for 95% Confidence Interval.

1.1.1 - Country: *

1.1.2 - EMCDDA data collection year *

1.1.3 - Year of the study: *

2. - Quantitative information: Lifetime prevalence

2.1. - EMCDDA Age range "All adults" (15-64)

2.1.1 - Age range used for "All adults" (if different from the EMCDDA) - LTP

2.1.2 - Prevalence (%) 15-64 - LTP

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
1. Any illegal drugs	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2. Cannabis (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
3. Opioids (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
4. Heroin (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
5. Other opioids (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
6. Cocaine (total, including crack) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
7. Amphetamines (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
8. Ecstasy (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
9. Hallucinogens (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
10. LSD (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
11. Other hallucinogens (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
12. Sedatives and/or tranquilisers (total) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
13. Barbiturates	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
14. Benzodiazepines	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
15. Solvents or inhalants	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
16. Anabolic steroids	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
17. Alcohol (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
18. Tobacco (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

2.2. - EMCDDA Age range "Young adults" (15-34)

2.2.1 - Age range used for "Young adults" (if different from the EMCDDA) - LTP

2.2.2 - Prevalence (%) 15-34 - LTP

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
1. Any illegal drugs	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2. Cannabis (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
3. Opioids (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
4. Heroin (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
5. Other opioids (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
6. Cocaine (total, including crack) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
7. Amphetamines (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
8. Ecstasy (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
9. Hallucinogens (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
10. LSD (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
11. Other hallucinogens (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
12. Sedatives and/or tranquilisers (total) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
13. Barbiturates	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
14. Benzodiazepines	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
15. Solvents or inhalants	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
16. Anabolic steroids	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
17. Alcohol (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
18. Tobacco (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

2.3. - Broad age groups

2.3.1 - EMCDDA Broad age groups (15-24) / (25-34) / (35-44) / (45-54) / (55-64) - LTP

	15-24	25-34	35-44	45-54	55-64
Age ranges used (if different from EMCDDA)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

2.3.2 - Prevalence (%) - Broad age groups LTP

	15-24 (M)	15-24 (F)	15-24 (total)	25-34 (M)	25-34 (F)	25-34 (total)	35-44 (M)	35-44 (F)	35-44 (total)	45-54 (M)	45-54 (F)	45-54 (total)	55-64 (M)	55-64 (F)	55-64 (total)
1. Any illegal drugs	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2. Cannabis (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
3. Opioids (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
4. Heroin (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
5. Other opioids (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
6. Cocaine (total, including crack)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

	15-24 (M)	15-24 (F)	15-24 (total)	25-34 (M)	25-34 (F)	25-34 (total)	35-44 (M)	35-44 (F)	35-44 (total)	45-54 (M)	45-54 (F)	45-54 (total)	55-64 (M)	55-64 (F)	55-64 (total)
(EMQ)															
7. Amphetamines (EMQ)															
8. Ecstasy (EMQ)															
9. Hallucinogens (total)															
10. LSD (EMQ)															
11. Other hallucinogens (specify)															
12. Sedatives and/or tranquilisers (total) (EMQ)															
13. Barbiturates															
14. Benzodiazepines															
15. Solvents or inhalants															
16. Anabolic steroids															
17. Alcohol (EMQ)															
18. Tobacco (EMQ)															

3. - Quantitative information: Last year prevalence

3.1. - EMCDDA Age range "All adults" (15-64)

3.1.1 - Age range used for "All adults" (if different from the EMCDDA) - LYP

3.1.2 - Prevalence (%) 15-64 - LYP

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
1. Any illegal drugs					
2. Cannabis (EMQ)					
3. Opioids (total)					
4. Heroin (EMQ)					
5. Other opioids (specify)					
6. Cocaine (total, including crack) (EMQ)					
7. Amphetamines (EMQ)					

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
8. Ecstasy (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
9. Hallucinogens (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
10. LSD (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
11. Other hallucinogens (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
12. Sedatives and/or tranquillisers (total) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
13. Barbiturates	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
14. Benzodiazepines	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
15. Solvents or inhalants	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
16. Anabolic steroids	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
17. Alcohol (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
18. Tobacco (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

3.2. - EMCDDA Age range "Young adults" (15-34)

3.2.1 - Age range used for "Young adults" (if different from the EMCDDA) - LYP

3.2.2 - Prevalence (%) 15-34 - LYP

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
1. Any illegal drugs	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2. Cannabis (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
3. Opioids (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
4. Heroin (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
5. Other opioids (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
6. Cocaine (total, including crack) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
7. Amphetamines (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
8. Ecstasy (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
9. Hallucinogens (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
10. LSD (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
11. Other hallucinogens (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
12. Sedatives and/or tranquillisers (total) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
13. Barbiturates	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
14. Benzodiazepines	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
15. Solvents or inhalants	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
16. Anabolic steroids	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
17. Alcohol (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
18. Tobacco (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

3.3. - Broad age groups

3.3.1 - EMCDDA Broad age groups (15-24) / (25-34) / (35-44) / (45-54) / (55-64) - LYP					
	15-24	25-34	35-44	45-54	55-64
Age ranges used (if different from EMCDDA)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

3.3.2 - Prevalence (%) - Broad age groups LYP															
	15-24 (M)	15-24 (F)	15-24 (total)	25-34 (M)	25-34 (F)	25-34 (total)	35-44 (M)	35-44 (F)	35-44 (total)	45-54 (M)	45-54 (F)	45-54 (total)	55-64 (M)	55-64 (F)	55-64 (total)
1. Any illegal drugs	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2. Cannabis (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
3. Opioids (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
4. Heroin (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
5. Other opioids (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
6. Cocaine (total, including crack) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
7. Amphetamines (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
8. Ecstasy (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
9. Hallucinogens (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
10. LSD (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
11. Other hallucinogens (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
12. Sedatives and/or tranquilisers (total) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
13. Barbiturates	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

	15-24 (M)	15-24 (F)	15-24 (total)	25-34 (M)	25-34 (F)	25-34 (total)	35-44 (M)	35-44 (F)	35-44 (total)	45-54 (M)	45-54 (F)	45-54 (total)	55-64 (M)	55-64 (F)	55-64 (total)
14. Benzodiazepines	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
15. Solvents or inhalants	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
16. Anabolic steroids	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
17. Alcohol (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
18. Tobacco (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

4. - Quantitative information: Last month prevalence

4.1. - EMCDDA Age range "All adults" (15-64)

4.1.1 - Age range used for "All adults" (if different from the EMCDDA) - LMP

4.1.2 - Prevalence (%) 15-64 - LMP

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
1. Any illegal drugs	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2. Cannabis (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
3. Opioids (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
4. Heroin (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
5. Other opioids (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
6. Cocaine (total, including crack) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
7. Amphetamines (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
8. Ecstasy (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
9. Hallucinogens (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
10. LSD (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
11. Other hallucinogens (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
12. Sedatives and/or tranquilisers (total) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
13. Barbiturates	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
14. Benzodiazepines	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
15. Solvents or inhalants	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
16. Anabolic steroids	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
17. Alcohol (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
18. Tobacco (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

4.2. - EMCDDA Age range "Young adults" (15-34)

4.2.1 - Age range used for "Young adults" (if different from the EMCDDA) - LMP

4.2.2 - Prevalence (%) 15-34 - LMP

	Male	Female	Total	Lower CI 95 (Total)	Upper CI 95 (Total)
1. Any illegal drugs	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2. Cannabis (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
3. Opioids (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
4. Heroin (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
5. Other opioids (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
6. Cocaine (total, including crack) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
7. Amphetamines (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
8. Ecstasy (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
9. Hallucinogens (total)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
10. LSD (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
11. Other hallucinogens (specify)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
12. Sedatives and/or tranquilisers (total) (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
13. Barbiturates	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
14. Benzodiazepines	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
15. Solvents or inhalants	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
16. Anabolic steroids	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
17. Alcohol (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
18. Tobacco (EMQ)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

4.3. - Broad age groups

4.3.1 - EMCDDA Broad age groups (15-24) / (25-34) / (35-44) / (45-54) / (55-64) - LMP

	15-24	25-34	35-44	45-54	55-64
Age ranges used (if different from	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

	15-24	25-34	35-44	45-54	55-64
EMCDDA)					

4.3.2 - Prevalence (%) - Broad age groups LMP

	15-24 (M)	15-24 (F)	15-24 (total)	25-34 (M)	25-34 (F)	25-34 (total)	35-44 (M)	35-44 (F)	35-44 (total)	45-54 (M)	45-54 (F)	45-54 (total)	55-64 (M)	55-64 (F)	55-64 (total)
1. Any illegal drugs															
2. Cannabis (EMQ)															
3. Opioids (total)															
4. Heroin (EMQ)															
5. Other opioids (specify)															
6. Cocaine (total, including crack) (EMQ)															
7. Amphetamines (EMQ)															
8. Ecstasy (EMQ)															
9. Hallucinogens (total)															
10. LSD (EMQ)															
11. Other hallucinogens (specify)															
12. Sedatives and/or tranquilisers (total) (EMQ)															
13. Barbiturates															
14. Benzodiazepines															
15. Solvents or inhalants															
16. Anabolic steroids															
17. Alcohol (EMQ)															
18. Tobacco (EMQ)															

5. - Survey details

5.1. - Methodology

5.1.1 - Reference (complete main bibliographic reference of the study)
--

5.1.2 - Survey website address

5.1.3 - Other publications (produced by the study)

5.1.4 - Year of the study:

5.1.5 - Institution commissioning (paying for) survey

5.1.6 - Research agency (if different from commissioning institution) (not the field agency)

5.1.7 - Single/repeated study

5.1.8 - Context (health/crime/drugs only)

5.1.9 - Geographical area covered*

5.1.10 - Total age range covered

5.1.11 - Mode of data collection (face-to-face, mail, phone,...)

5.1.12 - Sampling frame (population registry, household registry,...)

5.1.13 - Sampling procedures

5.1.14 - Oversampled groups

5.1.15 - Weighting procedures (by age, gender,...)

5.1.16 - Total gross sample size (-Nt-) (number)

(Nt) includes all cases initially selected, including wrong addresses, people where no contact could be made due to absence of respondent, etc

5.1.17 - Valid gross sample size (-Nv-) (number)

(Nv) similar to (Nt) but excluding frame errors (wrong or non-existing addresses, offices addresses...)

5.1.18 - Net response for "total age range covered" (-n-) (number)

(n) includes the number of interviews actually carried out. Often this is the figure referred as "sample size"

5.1.19 - Net response 15-64 years old (number)

5.1.20 - Net response 15-34 years old (number)

5.1.21 - Net response by age group

	15-24	25-34	35-44	45-54	55-64
Net response (15-24) / (25-34) / (35-44) / (45-54) / (55-64) (numbers)	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

5.1.22 - Response rates

	Male (%)	Female (%)	Total (%)
Response rates - total age range covered (percentage)	<input type="text"/>	<input type="text"/>	<input type="text"/>
Response rate 15-64 (percentage)	<input type="text"/>	<input type="text"/>	<input type="text"/>

	Male (%)	Female (%)	Total (%)
Response rate 15-34 (percentage)	<input type="text"/>	<input type="text"/>	<input type="text"/>

"Response rate" will be computed as $(n / Nt - \text{frame errors}) * 100$ -or the equivalent- $(n / Nv) * 100$. If computed in a different way, explain in question 5.2.23.

5.1.23 - Please describe in detail how response rates were computed in practice

5.1.24 - Remarks

5.1.25 - Drugs definitions

Provide a detailed description of what is included in each drug category

1. Any illegal drugs	<input type="text"/>
2. Cannabis (EMQ)	<input type="text"/>
3. Opioids (total)	<input type="text"/>
4. Heroin (EMQ)	<input type="text"/>
5. Other opioids (specify)	<input type="text"/>
6. Cocaine (total, including crack) (EMQ)	<input type="text"/>
7. Amphetamines (EMQ)	<input type="text"/>
8. Ecstasy (EMQ)	<input type="text"/>
9. Hallucinogens (total)	<input type="text"/>
10. LSD (EMQ)	<input type="text"/>
11. Other hallucinogens (specify)	<input type="text"/>
12. Sedatives and/or tranquilisers (total) (EMQ)	<input type="text"/>
13. Barbiturates	<input type="text"/>

Provide a detailed description of what is included in each drug category

14. Benzodiazepines	
15. Solvents or inhalants	
16. Anabolic steroids	
17. Alcohol (EMQ)	
18. Tobacco (EMQ)	